



THE LARGE SYNOPTIC SURVEY TELESCOPE



Ian Shipsey
(for the LSST Collaboration)

Birmingham
February 11, 2015


Progress in Astronomy

- Bigger Telescopes: *Keck to E-ELT*
- Angular resolution: *Hubble to JWST*
- All Sky Survey: *Sloan Digital Sky Survey to LSST*

LSST

wide fast deep

I. Shipsey



LSST

Enabled by Technology

**Sensors
Computing
Large Optics Fabrication**

LSST : an integrated survey system designed to conduct a decade-long, deep, wide, fast time-domain survey of the optical sky.


* 8-m class wide-field ground based telescope

* 3.2 Gpix camera

* automated data processing system

LSST in a nutshell

Synoptic =
Big Picture



Wide


Comparison of LSST To Keck

Primary mirror diameter


Field of view
(full moon is 0.5 degrees)




Keck Telescope




0.2 degrees



LSST



3.5 degrees




Outer Space - The Cosmos

The
Hubble
Deep
Field



Sun




Hubble deep field

**UNIVERSE
OF
GALAXIES**

3000
here



100 billion
over entire
sky




I. Shipsey

Image sizes LSST, Moon, HST




Shipsey

Survey Power = aperture x field of view




Survey Power = aperture x field of view



Fast

189 4K x 4K CCDs Largest & fast astronomy CCD camera




I. Shipsey

**3 Gpix
multiport CCDs**

**Record image in
15 seconds**

**Readout image
In 2 seconds**




Trends in Optical Astronomy Survey Data



DEEP




Kosmas IV c. a.C



SDSS XXI c. a.C.

DEEP

LSST Probes a Volume an Order of Magnitude Larger than Current or Near-Future Surveys



- LSST ~100 times fainter than the Sloan Digital Sky Survey
- a legacy dataset ~1000 times as large
- ~800 images of every field will open up the time domain for large-scale study for the first time

DEEP

LSST Probes a Volume an Order of Magnitude Larger than Current or Near-Future Surveys

- LSST ~100 times fainter than SDSS
- a legacy dataset ~1000 times as large
- ~800 images of every field will open up the time domain for large-scale study for the first time: a movie of the universe

Google Earth

Google[™]
Sky

Google Universe

DEEP

LSST Probes a Volume an Order of Magnitude Larger than Current or Near-Future Surveys

- LSST ~100 times fainter than SDSS
- a legacy dataset ~1000 times as large
- ~800 images of every field will open up the time domain for large-scale study for the first time: a movie of the universe

A survey of 40 billion objects in space and time

32 trillion measurements

4 billion galaxies with redshifts

Time domain:


5 million asteroids

10 million supernovae

1 million gravitational lenses

100 million variable stars

+ new phenomena



LSST survey of 18,000 sq deg (half the sky)

- **4 billion galaxies with redshifts**
- **Time domain:**
 - 5 million asteroids**
 - 10 million supernovae**
 - 1 million gravitational lenses**
 - 100 million variable stars**
 - + new phenomena**


LSST Wide-Fast-Deep survey

A survey of 40 billion objects
in space and time

32 trillion measurements


LSST 4 Science Missions

Dark Energy-Dark Matter



Multiple investigations into the nature of the dominant components of the universe

Inventory of the Solar System




Find 82% of hazardous NEOs down to 140 m over 10 yrs & test theories of solar system formation


“Movie” of the Universe: time domain



Discovering the transient & unknown on time scales days to years



Mapping the Milky Way




Map the rich and complex structure of the galaxy in unprecedented detail and extent

All missions conducted in parallel


LSST 4 Science Missions

Dark Energy-Dark Matter



Multiple investigations into the nature of the dominant components of the universe

Inventory of the Solar System



Find 82% of hazardous NEOs down to 140 m over 10 yrs & test theories of solar system formation


“Movie” of the Universe: time domain



Discovering the transient & unknown on time scales days to years



Mapping the Milky Way



Map the rich and complex structure of the galaxy in unprecedented detail and extent

All missions conducted in parallel (similar to a general purpose expt @ LHC)

Summary of High Level Requirements

Survey Property	Performance
Main Survey Area	18000 sq. deg.
Total visits per sky patch	825
Filter set	6 filters (ugrizy) from 320 to 1050nm
Single visit	2 x 15 second exposures
Single Visit Limiting Magnitude	u = 23.5; g = 24.8; r = 24.4; I = 23.9; z = 23.3; y = 22.1
Photometric calibration	2% absolute
Median delivered image quality	~ 0.7 arcsec. FWHM
Transient processing latency	60 sec after last visit exposure
Data release	Full reprocessing of survey data annually

The Science Opportunities are summarized in

Quick read:


LSST: FROM SCIENCE DRIVERS TO REFERENCE DESIGN AND ANTICIPATED DATA PRODUCTS

<http://arxiv.org/pdf/0805.2366>
(last update August 2014)

Reference:

<http://www.lsst.org/lsst/scibook>

Written by 11 science
collaborations




The idea: 1996

The need for a facility to survey the sky *Wide, Fast and Deep*, has been recognized for many years.

1996-2000 “*Dark Matter Telescope*”
Emphasized mapping dark matter

2000- “*LSST*”
Emphasized a broad range of science from the same multi-wavelength survey data

LSST has been highly ranked by numerous US Astronomy and Particle Physics Review committees
Including NRC Astronomy Decadal Survey: Astro2010



We have been going through the approval process @ DOE and NSF

NSF \$473M Telescope & Data Management
DOE \$168M Camera
Private \$40M (already received)
NSF \$270M operations (10 years)
Non-US \$100M operations (10 years)

Good Timing: The Green light



National Science Board of NSF authorised project construction Aug. 1 2014

For NSF, funding profile consistent with LSST proposed schedule.
Total NSF (telescope and data management) project cost of \$473M.

The DOE budget for the LSST Camera is also consistent with planned funding profile.: Estimated camera project cost \$168M.


Commissioning start 2020

Science October, 2022

US will provide 270M\$ for operations
non-US partners provide 100M\$ in return for data rights



Integrated Project Schedule with Key Milestones



LSST in the UK


A next generation wide field optical survey is well-justified by the cartography, cinematography and photometry it will perform and the huge range of astrophysics and physics at the boundary between particle physics and astrophysics it will address.

LSST is the missing piece in the UK's future ground-based astronomy programme

Astrophysicists at 33 UK institutions have recently formed LSST:UK and are seeking to join LSST as a national consortium

UK Involvement in the Large Synoptic Survey Telescope


Presented
to STFC
PPRP
October 27
Panel
follow-up
questions
and
site visit
Jan 9






The LSST : UK Consortium



LSST Science Drivers 1 The Fate of the Universe



Flat universe
 $\Omega_{\text{total}} = 1.02 \pm 0.02$
 WMAP+Planck




Dark Energy “the essence of space”
 Dark Matter “most of the matter”
 Together they govern the evolution & fate of the universe.

Nobel 2011

Their nature ranks as one of the greatest questions in the physical sciences

Evolution of the energy density of the universe : $\rho \propto a^{-3(1+w)}$



Dark Energy: An unprecedented opportunity



Either:


two thirds of the energy in the Universe is of unknown origin,

Or:


General Relativity is wrong at large scales

Challenge: determine origin of Dark Energy or disprove GR

Approach: measure DE equation of state, w and its evolution, to the systematic limit with *multiple probes*



Dark energy equation of state parameters:





Does DE density evolve?

YES



*Are DE observations
self-consistent within
general relativity?*




New form
of energy




New theory
of gravity

NO




Cosmological
constant

Studying Dark Energy is one of the ways we may bring one of the greatest prizes in Physics within reach: reconciliation of the two great edifices



General Relativity



Quantum Mechanics


Probing Dark Energy

- The observable probing the properties of dark energy is the expansion history of the universe, and parameterized by the Hubble parameter $H(z)$


$$H(z) = \frac{\dot{a}}{a}$$

- Cosmic distances are proportional to integrals of $H(z)^{-1}$ over redshift.
- $H(z)$ can be constrained by measuring

luminosity distances
of standard candles
(Type 1a SNe)




angular diameter
distances of
standard rulers
baryon acoustic
oscillations (BAO)




Probing Dark Energy

Measure the expansion history of the universe

**luminosity distances
of standard candles
(Type 1a SNe)**





**angular diameter
distances of
standard rulers
baryon acoustic
oscillations (BAO)**



**•measure growth of structure as
function of redshift**

**•Galaxy Cluster surveys & Weak
Lensing (WL) Surveys**




Probing Dark Energy

- Second approach: measure growth of structure as function of redshift

- Stars, galaxies, clusters of galaxies grow by gravitational instability as the universe cools.


- Acceleration: The stretching of space – shuts off growth by keeping galaxies apart

- A cosmic “clock”




Measuring growth history, i.e. - the redshift at which structures of a given mass start to form is sensitive to the level of acceleration i.e. amount of dark energy

- Galaxy Cluster surveys & Weak Lensing (WL) Surveys probe growth of structure as well as angular diameter distances




Gravitational Lensing



$$\alpha = 4 G M / (c^2 b)$$

Bending of star light (point-like) by the Sun - star appears displaced



Newton 1704

Soldner 1801

$$\theta_N = \frac{2GM}{rc^2}$$

Einstein 1915


$$\theta_E = \frac{4GM}{rc^2}$$

Eddington 1919 → “between 1.59 and 1.86 arc-sec”





Extended objects are sheared

Red galaxy on axis strongly lensed. other galaxies weakly lensed: sheared images





Circular bkgd galaxies



what is observed

variable shape bkgd galaxies



Weak Lensing shear pattern less obvious but detectable statistically


Dark matter replaces the sun & distant galaxies replace the stars

Strong lensing requires alignment, rare, readily visible

Weak lensing, does not require alignment, common, detectable only statistically

Cosmic Shear

- **Cosmic Shear is the systematic and correlated distortion of the appearance of background galaxies due to weak gravitational lensing by the clustering of dark matter in the intervening universe.**
- **A given galaxy image is both displaced and sheared.**
- **The effect is detectable only statistically. *The shearing of neighboring galaxies is correlated, because their light follows similar paths on the way to earth.***



Tyson et al 2002

- **Massively exaggerated**

Cosmic shear: ~ 0.01


e.g. circular galaxy \rightarrow ellipse with $a/b \sim 1.01$

1st Detections of Cosmic Shear (2000-2003)

The simplest measure of cosmic shear is the 2-pt correlation function of the ellipticities measured with respect to angular scale.

Whitman 2000
145,000 galaxies
~1 degree

$\langle e(r) \bullet e(r + \theta) \rangle$
Log ellipticity correlation



No dark energy
Means more structure & more shear

$\Omega(\text{DE}) = 0.67$

Universe at $\frac{1}{2}$ critical density

Green dash galaxies are x2 closer


More recent surveys
CFHTLenS (2013)
4.2 million galaxies
~154 sq degree

SDSS (2011)
4.7 million galaxies
~275 sq degree

Lensing tomography

As statistics grow measurement of cosmic shear as a function of redshift becomes possible

Galaxies at z_2 will experience more shear than z_1



CFHTLenS(2013)
4.2 million galaxies
~154 sq degree


LSST and Cosmic Shear

CFHTLenS (2013)
4.2 million galaxies
~154 sq degree




LSST
4 billion galaxies
18,000 sq. degrees

- Same 2-pt correlation function
- Fourier transform \rightarrow power spectrum as a function of multi-pole moment (similar to CMB temperature maps).
- The growth in the shear power spectrum with the red shift of the background galaxies is provides the constraints on dark energy.






SDSS (2011)
4.7E6 galaxies
~275 sq degree



Baryon Acoustic Oscillations

- Prior to the formation of atoms baryons are tightly coupled to the radiation in the universe.
- An overdensity perturbation gives rise to an acoustic wave in this tightly coupled fluid, which propagates outward at the sound speed,
- After recombination, the matter and radiation decouple. The sound speed drops to zero, and the propagating acoustic wave stops.
- This gives rise to a characteristic scale in the universe: 150 Mpc the distance the sound waves have traveled at the time of recombination.




These acoustic waves are visible as the peaks in the CMB power spectrum.

Baryon Acoustic Oscillations

- Following recombination, gravitational instability causes the birth of stars and galaxies.
- Gravitational coupling between dark matter and baryons creates an imprint of the acoustic oscillations in the galaxy distribution.
- This persists as the universe expands, although it gets weaker with time.

$r_s \sim 150 \text{ Mpc}$






1.6°
at $z=2$

0.5°
moon


Baryon Acoustic Oscillations

$r_s \sim 150 \text{ Mpc}$



Baryon Acoustic Oscillations



$r_s \sim 150 \text{ Mpc}$





linear superposition

Same physics as CMB ($Z \sim 1100$)
but at a time when Dark
Energy is becoming important ($z < 3$)

Baryon Acoustic Oscillations SDSS




 1st observation
SDSS
Eisenstein
et al (2005)
40,000 galaxies
 $0.16 < z < 0.47$





1st observation
SDSS Eisenstein
et al (2005)
40,000 galaxies
 $0.16 < z < 0.47$



BOSS (2013)
1 million galaxies
 $8,500 \text{ deg}^2$ 13 Gpc^3
largest survey to date
 $z=0.32$ & $z=0.57$




Measure angle θ , trigonometry to get distance



I. Shipsey

- How the length scale evolves with redshift is sensitive to the expansion history of the universe and therefore to dark energy



BOSS (2013)
1 million galaxies






LSST
3 billion galaxies

WiggleZ Katzin 2014
BOSS Anderson 2013
6dFGS Beutler 2011

Baryon Acoustic Oscillations and LSST

Compilation of data this time as a power spectrum





I. Shipsey

BOSS (2013)
1 million galaxies



LSST
3 billion galaxies

Simulations of LSST measured galaxy power spectrum divided by a featureless reference power spectrum, shifted vertically for clarity



3. Supernovae



- Roughly 10^3 supernovae have been discovered to date
- LSST will find $> 10^7$ over its ten-year duration, spanning a broad redshift range, with precise, uniform calibration.
- This will revolutionize the field, allowing large samples for studies of systematic effects and additional parametric dependences.
- $\sim 10^5$ SNe Ia will be found in the “deep drilling fields” with well-measured lightcurves in all six colors. This will be an excellent sample for precision cosmology.
- The large sample size will also allow for the first time to conduct SN Ia cosmology experiments as a function of direction in the sky, providing stringent tests of the fundamental cosmological assumptions of homogeneity and isotropy.

I. Shipsey


3. Supernovae



- Roughly 10^3 supernovae have been discovered to date
- LSST will find $> 10^7$ over its ten-year duration, spanning a broad redshift range, with precise, uniform calibration.
- This will revolutionize the field, allowing large samples for studies of systematic effects and additional parametric dependences.
- The large sample size will also allow for the first time to conduct SN Ia cosmology experiments as a function of direction in the sky, providing stringent tests of the fundamental cosmological assumptions of homogeneity and isotropy.

I. Shipsey


Isotropy of Cosmic Acceleration



Current constraints on Dark Energy from multiple techniques

$$w = P / \rho$$
$$w = w_0 + w_a \left(\frac{z}{1+z} \right)$$

now **evolution**



Planck
arXiv:1303.5076v2
December, 2013

Combined:
SN + BAO + CMB


Predicted LSST Constraints on Dark Energy from multiple techniques

$$w = P / \rho$$


$$w = w_0 + w_a \left(\frac{z}{1+z} \right)$$

now

evolution



→
Present state
of knowledge



Science Driver 2: Mapping the Milky Way

An SDSS image of the Cygnus Region

With LSST:

About 200 images, each 2 mag deeper

The co-added images will be 5 mag. deeper

Precise proper motion & parallax measurements
will be available for $r < 24$ (4 magnitudes deeper
than the Gaia survey)

Stellar Populations



- **LSST will individually resolve and detect billions of stars in the Milky Way and neighboring Local Group galaxies,**
- **Studies of field stars and stellar associations can address a multitude of astrophysical issues associated with star formation and evolution, the assembly of the MW galaxy, and the origin of the chemical elements.**
- **Key techniques for these investigations include:**
 - **Construction of color magnitude diagrams**
 - **Trigonometric parallaxes** to establish absolute distances
 - **Stellar proper motions** to separate associations from background stars and from one another
 - Using **RR Lyrae** and other variables as “standard candles”
 - Using **eclipsing binaries** to measure stellar masses


Example: structure of outer milky way

RR Lyrae stars are luminous enough and copious enough to map the outer galaxy

Overdensities found in SDSS star count studies to 100 kpc

LSST RR Lyrae to 400 kpc, extending SDSS mapping volume by a factor of 50.

An important test of the small-scale accretion history of the Galaxy and a test of standard Model of cosmology



The standard model of cosmology predicts that the Milky Way should have accreted and destroyed hundreds of small dwarf galaxies in the past 10 Gyr. The residue survives as structure (star over-densities) in the outer halo.

Image: Star density stellar halo simulations

Science Driver 3 Inventory of the Solar System

Example: Near Earth Objects

- Inventory of solar system is incomplete
Estimate 17,000 undetected
- LSST would determine orbits of nearly all NEOs larger than 150m
- Demanding project: requires mapping the sky down to 24th magnitude every few days, individual exposures not to exceed 15 sec

The Sky is Falling


- Meteoroids/Fireballs that are golf-ball sized and up
 - Each day, ~ 100 tons of rock burns up in our atmosphere.


- **This fireball witnessed by thousands of people on October 9, 1992 in... guess where?**
 - streaked across sky at 50,000 km/h
 - 1st meteor ever filmed *and then* recovered





Killer Meteorites?







Potentially Hazardous Asteroids

4000 estimated

600 charted




I. Shipsey



Tunguska
(1908)
40 m asteroid

Recent Close Calls

Many near misses are discovered only after the object has missed Earth.



■ March 18, 2004


- 100 m asteroid came within 43,000 km of Earth
- discovered 1 day in advance

■ Similar events happen roughly every

- 500 m diameter asteroid comes within 100,000 km of Earth

October 27 2014 2014 UF56 0.4 x lunar distance
June 3 2014 2014 LY21 0.02 x lunar distance
Feb 8 2015 2013 CL 4.3 x lunar distance
<http://neo.jpl.nasa.gov/ca/>


Percentage of Potentially Hazardous Asteroids Found



Understanding the formation and evolution of the Solar System

- LSST will detect and determine orbits for millions of small bodies in the Solar System.
- Classes include:
 - [Near Earth Asteroids \(NEAs\)](#), and their subclass, [Potentially Hazardous Asteroids \(PHAs\)](#), whose orbits can potentially impact the Earth.
 - [Main Belt Asteroids \(MBAs\)](#), lying between the orbits of Mars and Jupiter.
 - [Trojans](#), which are asteroids in 1:1 mean motion resonance with a planet.
 - [Trans-Neptunian Objects \(TNOs\)](#), and their subclass, [Classical Kuiper Belt Objects \(cKBOs\)](#). These occupy a large area of stable orbital space.
 - [Jupiter-Family Comets \(JFCs\)](#), whose orbits are strongly perturbed by Jupiter.
 - [Long Period Comets \(LPCs\)](#), which originate in the Oort Cloud at 10,000 AU.
 - [Halley Family Comets \(HFCs\)](#), which also come from the Oort Cloud, but have shorter periods.
 - [Damocloids](#), a group of asteroids with similar dynamical properties to the HFCs.
- Understanding the origin and behavior of these various systems is crucial for modelling the formation and evolution of the Solar System.

Science Driver 4: Transients & variable objects



Supernova

Optical
flashes

Optical
flashes

Deep Lens Survey

Becker, A. C., et al. 2004, Astrophysical Journal, 611, 418
f. Shipsey

Science Driver 4: Transients and Variable Objects

Recent surveys have shown the power of measuring variability for studying gravitational lensing, searching for supernovae, determining the physical properties of gamma-ray burst sources, probing the structure of active galactic nuclei, studying variable stars, and many other subjects at the forefront of astrophysics.

Wide-area, dense temporal coverage to deep limiting magnitudes enables the discovery and analysis of rare and exotic objects such as neutron star and black hole binaries, novae and stellar flares, gamma-ray bursts and X-ray flashes, active galactic nuclei (AGNs), stellar disruptions by black holes, and possibly new classes of transients, such as binary mergers of black holes.


LSST: ~10 million cosmic explosions over most of the observable Universe, extending the volume of the parameter space for discovery by $\times 1,000$ reaching unprecedented sensitivity. A movie of the universe



Massively Parallel Astrophysics

- **Dark matter/dark energy via weak lensing**
- **Dark energy via baryon acoustic oscillations**
- **Dark energy via supernovae**
- **Galactic Structure encompassing local group**
- **Dense astrometry over 18000 sq.deg: rare moving objects**
- **Gamma Ray Bursts and transients to high redshift**
- **Gravitational micro-lensing**
- **Strong galaxy & cluster lensing: physics of dark matter**
- **Multi-image lensed SN time delays: separate test of cosmology**
- **Variable stars/galaxies: black hole accretion**
- **Optical bursters to 25 mag: the unknown**
- **5-band 27 mag photometric survey: unprecedented volume**
- **Solar System Probes: Earth-crossing asteroids, Comets, TNOs**
- **Planetary transits**

All science programs conducted in parallel




LSST Optical Design


I. Shipsey

LSST Optical Design

- $f/1.23$ Very short focal length gives wide field of view for given image size
- 3.5° FOV over a 64 cm focal plane, Etendue = $319 \text{ m}^2\text{deg}^2$
- < 0.20 arcsec FWHM images in six filter bands: $0.3 - 1 \mu\text{m}$




Cross section through telescope and camera



I. Shipsey

The primary/tertiary mirror is a long lead time item...

Stewart Observatory Mirror
Lab Tucson, AZ



High Fire, March 29 2008


1165°C (2125°F). Then anneal & cool gradually to room temp.

Mirror has been ground, and polished

Completion :2015




2 September 2008



I. Shipsey

Unique Monolithic M_1 / M_3 mirror polishing nearly complete



March 2014



Secondary Mirror Substrate ready for optical polishing

CHARLES AND LISA SIMONYI FUND
... FOR ARTS AND SCIENCES ...



March 2008



Sept 2008

Private donors have joined with more than thirty universities, research institutions, and corporations to construct the Large Synoptic Survey Telescope (LSST), a survey telescope capable of producing a high-definition digital movie of the entire sky and forever changing the way we view the Universe. The LSST Corporation invites you and a guest to share with us in the completion milestone of this unique telescope mirror, spun cast from molten glass in 2008 and meticulously polished to perfection for more than 6 years.

Save the Date

M1/M3 Completion

Saturday, January 10, 2015


Please join us on January 10th, 2015, to celebrate the completion of the monolithic primary (M1) and tertiary (M3) mirrors for the LSST at the University of Arizona's Steward Observatory Mirror Lab in Tucson, Arizona.

A formal invitation with details regarding scheduled events, hotel, and more will follow. For now, just save the date!

Questions can be directed to:



Pat Eliason, peliason@lsst.org | Sandra Ortiz, sortiz@lsst.org

More information about the Large Synoptic Survey Telescope can be found at <http://www.lsst.org>



I. Ships





M2 final polishing started
October 2014

THE POWER OF INGENUITY

LSST Will be Sited in Central Chile






LSST is
located in
a NSF
compound
near SOAR
& Gemini

I. Shipsey

Cerro Pachón, as
seen from Tololo,
April 9, 2011
(During first ever
LSST Board
meeting in Chile)



Site and observatory



facility designed to minimize atmospheric turbulence in the vicinity of the dome
Formal “laying of the 1st stone” for the observatory April 14, 2015

After ~4,000 kg of explosives and ~12,500 m³ of rock removal, Stage I of the El Peñón summit leveling is completed.



I. Shipsey



87

LSST Observing Cadence Set by Science Goals

Pairs of 15 second exposures (*to 24.5 mag*) per visit to a given position in the sky.

Visit the same position again within the hour with another pair of exposures.


Number of 9.6 sq.deg field-of-view visits per night: 900

**Detection of transients announced worldwide within
60 seconds.**

Expect 1-2 million alerts per night!


Telescope System Designed to Slew and Settle within 5 seconds

- The high curvature mirrors allow a short, light, stiff, stable and agile telescope employing an alt-azimuth mount
- Points to new positions in the sky every 39 seconds
- Tracks during exposures and slews 3.5° to adjacent fields in ~ 4 seconds
 - Moving Structure 350 tons (60 tons optical systems).
 - Pier design structured to maximize stiffness.





Telescope model with system design details included

FEA model is loaded structure on bearings, pier, and summit rock



Optical Quality at the LSST site





SDSS Apache Point NM, 1.3 arc sec seeing

Optical Quality at the LSST site

These two images are of the same patch of sky

What a telescope sees in Chile ,
very close to the LSST site
0.67 arcsec seeing




SDSS Apache Point NM, 1.3 arc sec seeing

x2 better seeing x5 fainter per image

(1,000 images at each sky location
will be obtained over 10 years, the
Coaddition is x75 fainter than SDSS)

....and for a single galaxy

SDSS




....and for a single galaxy

These two images are of the same galaxy




MUSYC



MUSYC is x25 fainter than SDSS but still x
X3 less faint than LSST

Gawiser et al


3.2 Gigapixel Camera



LSST's Six Optical Filter Bands


Determine color and redshift

Transmission- atmosphere, telescope, & detector QE




→ Photometric determination of galaxy redshifts

Photometric Redshifts



- **Galaxies have distinct spectra, with characteristic features at known rest wavelengths.**
- **Accurate redshifts can be obtained by taking spectra of each galaxy. But this is impractical for the billions of galaxies in LSST cosmic shear and BAO studies.**
- **Instead, the colors of the galaxies are obtained from the images themselves. This requires accurate calibration of both the photometry and of the intrinsic galaxy spectra as a function of redshift.**

LSST Camera: 21 science rafts, 189 4K x 4K CCDs



LSST prototype sensors meet project requirements.


1st article sensor procurement under way with two vendors e2v and ITL

Sensor delivery rate is the critical path pacing item for the LSST camera.


- Every 15 sec exposure, 2 sec readout, repeat
- 5 second slew to new sky location
- Nightly data generation rate: 15 Tbytes
- SDSS Data Release 7 was 16TB

LSST Camera:

21 science rafts, 189 4K x 4K CCDs





Science Raft Comprises 9 CCDs and associated electronics.




Individual sensor includes mechanical mount and flex cable
Each raft is a standalone 144 Mpix camera

Focal Plane Sensors Quantum Efficiency





Quantum Efficiency


Vendor Data $t=100\ \mu\text{m}$




LSST (BNL) DATA




The LSST sensors have been tested on-the sky



Vendor 2 buttable package



Vendor 1 4K x 4K device in test Dewar



**also known as Calypso at Kitt Peak*

Sensors meet Requirements, Procurement is Under Way

LSST prototype sensors meet project requirements.

1st article sensor procurement is now under way with two vendors e2v and ITL

Sensor delivery rate is the critical path pacing item for the LSST camera.

prototype, vendor 1



prototype, vendor 2




compliance matrix

	Number	Specification	Value	Unit	No. Tested/Passes	Result
ARCHITECTURE	CCD-001	Format	Design Consideration		ALL	-
	CCD-002	Pixel size	Design Consideration		ALL	-
	CCD-003	Segmentation	Design Consideration		ALL	-
	CCD-004	Contiguity	Design Consideration		ALL	-
ELECTRICAL	CCD-006	Read time	2	sec	9/9	tested at 545kpix/s
	CCD-007	Read noise	8	e- rms	6/0	7.8 ±1
					3/3	5.01 ± 0.97
	CCD-008	Bloomed full well	175000	e- max	4/4	145000 ± 17000
	CCD-009	Nonlinearity	±2	%	2/2	1.00 ± 0.3

- Every 15 sec exposure, 2 sec readout, repeat
- 5 second slew to new sky location
- Nightly data generation rate: 15 Tbytes
- SDSS Data Release 7 was 16TB

LSST Operations Sites and Data Flows



Archive Site

Archive Center

Alert Production
Data Release Production
Calibration Products Production
EPO Infrastructure
Long-term Storage (copy 2)


Data Access Center

Data Access and User Services

Dedicated Long Haul Networks


Two redundant 40 Gbit links from La Serena to Champaign, IL (existing fiber)

Summit and Base Sites




Telescope and Camera
Data Acquisition
Crosstalk Correction
Long-term storage (copy 1)
Chilean Data Access Center

HQ Site



Science Operations
Observatory Management
Education and Public Outreach

Data volumes & rates are unprecedented in astronomy



Ultimate LSST Deliverable: Reduced Data Products



A petascale supercomputing system at the **LSST Archive** (at NCSA) will process the raw data, generating reduced image products, time-domain alerts, and catalogs.

Large Synoptic Survey Telescope
The widest, fastest, deepest eye of the new digital age

Searches History Read FITS File Preferences Catalogs Plot Layers Background Monitor

Search by Position 21.41;0.13;EQ_J2000; Type-CENTER; Filter=all; Image Size=0.0278 deg

LSST Image Data

goodSeeingCoaddId	tract	patch	filterName	ra	dec	fluxMag0	fluxMag2Sigma	measuredWhm
19922944	0	304,0	u	21.458185000	0.104445058	6.20437012e+10	0.000000	1.699982
19922945	0	304,0	g	21.458185000	0.104445058	6.22980014e+10	0.000000	1.699982
19922946	0	304,0	r	21.458185000	0.104445058	6.43898982e+10	0.000000	1.699982
19922947	0	304,0	i	21.458185000	0.104445058	6.58835005e+10	0.000000	1.699982
19922948	0	304,0	z	21.458185000	0.104445058	6.12743987e+10	0.000000	1.699982

LSST Multi-Color 1.2x

LSST Filter u 1x LSST Filter g 1x LSST Filter r 1x LSST Filter i 1x LSST Filter z 1x

Data Access Centers in the U.S. and Chile will provide end-user analysis capabilities and serve the data products to LSST users.

LSST From the User's Perspective

- Images
- A stream of ~10 million time-domain events per night, detected and transmitted to event distribution networks within 60 seconds of observation.
- A catalog of orbits for ~6 million bodies in the Solar System.

- A catalog of ~40 billion objects (20B galaxies, 20B stars), produced annually, accessible through online databases.
- Deep co-added images.

- Services and computing resources at the Data Access Centers to enable user-specified custom processing and analysis.
- Software and Applications Programming Interfaces enabling development of analysis codes.

Level 1
Nightly

Level 2
Annual

Level 3


Adaptive Optics and Active Optics

$$E = Ae^{i\phi}$$



Adaptive Optics (Rapid 50-300Hz)
 Limited to a small field of view
 (LSST has a big field of view)

Active Optics (LSST)

- * Measure perturbed wavefront to correct distortions in telescope and camera optics
- * BUT Long-exposure sampling of wavefront to average atmospheric turbulence
- * Telescope optical surfaces are adjustable between exposures to correct for distortions but remain static during each exposure




LSST Wavefront Sensing




Purdue/NOAO


Wavefront Sensing Corner Rafts of LSST Camera



CCD Curvature Sensor



Purdue/NOAO




Corner Raft mechanical & thermal design work


- Design for accurate and stable mount for sensors and electronics in the Camera
- Assembly sequence & insertion tooling
- Mechanical & Thermal analysis (FEA & prototype tests)
- risk & cost analysis

Purdue


Overall FE model $\Delta T = 30C$







ΔT across Corner Raft + sensors = 7.5C




Corner Raft surface deformation <1 micron



Wavefront Reconstruction and Sensor Evaluation Station at Purdue



Pressure and temperature curve LN2 + turbo + ion pumps



RGA scan

Status – all operational

- Cryostat (LN2 cooling + vacuum system)
- X-ray (Fe55) source
- Optical flat-field source
- Monochromator
- Electronic shutter
- Camera lens + motion control



Purdue

Initially configured for tests using single sensors for wavefront and guider studies.

The test station will be expanded to accommodate tests of a full Corner Raft/ Tower which will be fabricated @ Purdue

I. Shipsey


TESTBENCH FOR WAVEFRONT SENSING



Purdue


When 2um of x-axis astigmatism is dialed in to simulate a 2 micron distortion of the LSST primary mirror

Example



Intra-focal image
Z = -1mm

Extra-focal image
Z = +1mm



The two CCD images are used to Reconstruct the perturbed Wavefront . Fit it to get→





Image Simulation: Implementing a simulated sky






Image Simulation: Implementing a simulated sky



+ operations simulator

Following the photon flow...



https://dl.dropboxusercontent.com/u/24655052/movie_peterson.mpeg

Optics



+Tracking



+Diffraction



+Detector
Misalignments &
Perturbations



+Lens Misalignments



+Mirror Misalignments
Perturbations,
& Micro-roughness



+Detector



+High Altitude
Atmosphere



+Mid Altitude
Atmosphere



+Low Altitude
Atmosphere



+Pixelization




+Saturation &
Blooming



20 x 20 pixels

Simulated L (one exposure)


Three filter (gri)
composite
image 1CCD



Simulation of full LSST focal plane



**Simulation at the
scale of LSST
with the same
cadence
& similar
Systematics
Is a powerful
Probe of
Physics reach
& survey design**



LSST operations simulator

LSST Operations are determined by a special simulation program including real weather data, seeing, twilight, sky background (lunar), time to slew, overall survey coverage + and depth already achieved → ranking algorithm for next observation (constantly updated) results in the visits per patch of sky (color coded at right) for each of the six filters for 10 year survey at right






Figure : Visits numbers per field for the 10 year simulated survey


Science
Driver 1
Dark
Energy

→
Present state
of knowledge



The dark energy facilities roadmap





Measurements of dark energy with several techniques with Stage III experiments. Precision is limited.

Stage III

LSST
& other stage IV experiments

LSST Outreach Data will be used in classrooms, science museums, and online



Classroom Emphasis on:

- Data-enabled research experiences
- Citizen Science
- College classes
- Collaboration through Social Networking



LSST Education & Public Outreach

- **LSST is Telescope for Everyone**

LSST will discover 10 billion new galaxies— enough for everyone

Reaching for the sky has always inspired the deepest questions and boldest expeditions of discovery.

Now we can reach more of the Universe, through the vastness of time, in unprecedented detail.

A school child in South Africa, Chile, or Birmingham can discover an island universe


LSST Institutions



- The University of Arizona
- University of Washington
- National Optical Astronomy Observatory
- Research Corporation for Science Advancement
- Adler Planetarium
- Brookhaven National Laboratory (BNL)
- California Institute of Technology
- Carnegie Mellon University
- Chile
- Cornell University
- Drexel University
- Fermi National Accelerator Laboratory
- George Mason University
- Google, Inc.
- Harvard-Smithsonian Center for Astrophysics
- Institut de Physique Nucléaire et de Physique des Particules (IN2P3)
- Johns Hopkins University
- Kavli Institute for Particle Astrophysics and Cosmology (KIPAC) - Stanford University
- Las Cumbres Observatory Global Telescope Network, Inc.
- Lawrence Livermore National Laboratory (LLNL)
- Los Alamos National Laboratory (LANL)
- Northwestern University
- Princeton University
- Purdue University
- Rutgers University
- SLAC National Accelerator Laboratory
- Space Telescope Science Institute
- Texas A & M University
- The Pennsylvania State University
- University of California at Davis
- University of California at Irvine
- University of Illinois at Urbana-Champaign
- University of Michigan
- University of Oxford
- University of Pennsylvania
- University of Pittsburgh
- Vanderbilt University

...LSST is growing other UK groups are in the process of joining as are many others from around the globe

LSST IS HIRING



WE'RE SEEKING TOP TALENT TO WORK IN A TEAM ENVIRONMENT THAT INSPIRES EXCELLENCE.



JOIN US IN:

LSST HEADQUARTERS
TUCSON, AZ

SLAC/STANFORD
MENLO PARK, CA

PRINCETON UNIVERSITY
PRINCETON, NJ

NCSA / UIUC
URBANA-CHAMPAIGN, IL

UNIVERSITY OF WASHINGTON
SEATTLE, WA

LSST OBSERVATORY SITE
CERRO PACHÓN, CHILE

ABOUT US

LSST IS A PUBLIC-PRIVATE PARTNERSHIP AND THE TOP-RANKED LARGE-SCALE GROUND-BASED PROJECT FOR THE NEXT DECADE AS RECOMMENDED BY THE NRC'S ASTRO2010 DECADAL SURVEY. LSST WILL SCAN THE SKY FOR 10 YEARS, PRODUCING A PETABYTE-SCALE, NON-PROPRIETARY DATABASE DESIGNED TO ADDRESS THE MOST PRESSING QUESTIONS IN ASTRONOMY AND PHYSICS, WHILE DRIVING ADVANCES IN BIG-DATA SCIENCE AND COMPUTING.

CAREERS

LSST IS A NEW PARADIGM FOR LARGE SCIENTIFIC FACILITIES: OPEN SOURCE, OPEN DATA, AND AN OPEN, FLEXIBLE WORK ENVIRONMENT. ALL LSST WORK SITES OFFER EXCEPTIONAL BENEFITS PACKAGES AND ROOM FOR PROFESSIONAL GROWTH.

OUR TEAM

PROJECT OFFICE
DATA MANAGEMENT
TELESCOPE & SITE
CAMERA
EDUCATION & PUBLIC OUTREACH
SYSTEM ENGINEERING



 @LSST @mjuric

[LSST.ORG/HIRING](https://lsst.org/hiring)



CHARLES AND LISA SIMONYI FUND
••• FOR ARTS AND SCIENCES •••


Part of the LSST Collaboration 8/2012



**A partnership of particle physicists,
astrophysicists & computer scientists**

I. Shipsey

Summary



- **The Project Team is ready for a construction start in July 2014 to build the system to survey, store, process and serve the data starting in 2022**

Summary of the LSST Project

- The Project Team began construction in August 2014. We are building the system to survey, store, process and serve the data starting in 2022.
- The LSST science opportunities are extremely rich - ranging from studies of the smallest objects in the solar system to the structure and dynamics of the Universe as a whole.
- Most of the requisite investigations can be performed using data from a single coherent survey program. This is “massively parallel survey astrophysics” in its purest form.
- The analyses will be complex and will require significant attention to detailed systematics uncertainties. *There are many opportunities for astronomers, particle physicists and computer scientists to become involved now in helping us to optimize the anticipated science that will come from this marvelous facility.*