



H2020 MSCA COFUND  
G.A. 754496

# Stefano Gariazzo

*INFN, Turin section  
Turin (IT)*



Istituto Nazionale di Fisica Nucleare  
SEZIONE DI TORINO

[gariazzo@to.infn.it](mailto:gariazzo@to.infn.it)

<http://personalpages.to.infn.it/~gariazzo/>

# (Cosmological) Relic neutrinos, from A to Z

Seminar at the University of Birmingham, online, 03/03/2021


# A

# Active neutrinos


Spoiler: “Sterile” will come later

Based on:

- Planck 2018
- arxiv:2012.02726




# The Standard Model of Particle Physics



[D. Galbraith & C. Burgard, 2012]

# The Standard Model of Particle Physics




[D. Galbraith & C. Burgard, 2012]

# Neutrino oscillations


Major discoveries:

[SuperKamiokande, 1998]



first discovery of  $\nu_\mu \rightarrow \nu_\tau$   
oscillations from atmospheric  $\nu$

[SNO, 2001-2002]



first discovery of  $\nu_e \rightarrow \nu_\mu, \nu_\tau$   
oscillations from solar  $\nu$

Nobel prize in 2015


## Two neutrino bases

flavor neutrinos  $\nu_\alpha$

$$|\nu_\alpha\rangle = \sum_k U_{\alpha k} |\nu_k\rangle$$

massive neutrinos  $\nu_k$

$$|\nu(t=0)\rangle = |\nu_\alpha\rangle = U_{\alpha 1} |\nu_1\rangle + U_{\alpha 2} |\nu_2\rangle + U_{\alpha 3} |\nu_3\rangle$$



$$|\nu(t > 0)\rangle = |\nu_\beta\rangle = U_{\alpha 1} e^{-iE_1 t} |\nu_1\rangle + U_{\alpha 2} e^{-iE_2 t} |\nu_2\rangle + U_{\alpha 3} e^{-iE_3 t} |\nu_3\rangle \neq |\nu_\alpha\rangle$$

$$E_k^2 = p^2 + m_k^2 \xleftarrow{\text{define}} t = L$$

$$P_{\nu_\alpha \rightarrow \nu_\beta}(L) = |\langle \nu_\alpha | \nu(L) \rangle|^2 = \sum_{k,j} U_{\beta k} U_{\alpha k}^* U_{\beta j}^* U_{\alpha j} \exp\left(-i \frac{\Delta m_{kj}^2 L}{2E}\right)$$

$$\Delta m_{ij}^2 = m_i^2 - m_j^2$$

## The mixing matrix

$U$  can be parameterized using 3 angles ( $\theta_{12}$ ,  $\theta_{13}$ ,  $\theta_{23}$ ) and max 3 (1 Dirac  $\delta$ , 2 Majorana [ $\exists$  only for Majorana  $\nu$ ]) phases

$$U = \underbrace{\begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 0 & -s_{23} & c_{23} \end{pmatrix}}_{\text{mainly atmospheric and LBL accelerator disappearance}} \underbrace{\begin{pmatrix} c_{13} & 0 & s_{13} e^{-i\delta} \\ 0 & 1 & 0 \\ -s_{13} e^{i\delta} & 0 & c_{13} \end{pmatrix}}_{\text{mainly SBL reactors and LBL accelerator appearance}} \underbrace{\begin{pmatrix} c_{12} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix}}_{\text{mainly solar and LBL reactors}} M$$

Majorana phases irrelevant for oscillation experiments

Relevant for example in neutrinoless double-beta decay

$$s_{ij} \equiv \sin \theta_{ij}; \quad c_{ij} \equiv \cos \theta_{ij}$$

SBL = short baseline; LBL = long baseline

# Three Neutrino Oscillations

$$\nu_\alpha = \sum_{k=1}^3 U_{\alpha k} \nu_k \quad (\alpha = e, \mu, \tau)$$

$U_{\alpha k}$  described by 3 mixing angles  $\theta_{12}$ ,  $\theta_{13}$ ,  $\theta_{23}$  and one CP phase  $\delta$


Current knowledge of the 3 active  $\nu$  mixing: [JHEP 02 (2021)]

NO/NH: Normal Ordering/Hierarchy,  $m_1 < m_2 < m_3$

IO/IH: Inverted O/H,  $m_3 < m_1 < m_2$


$$\begin{aligned}\Delta m_{21}^2 &= (7.50^{+0.22}_{-0.20}) \cdot 10^{-5} \text{ eV}^2 \\ |\Delta m_{31}| &= (2.56^{+0.03}_{-0.04}) \cdot 10^{-3} \text{ eV}^2 \text{ (NO)} \\ &= (2.46 \pm 0.03) \cdot 10^{-3} \text{ eV}^2 \text{ (IO)}\end{aligned}$$

$$\begin{aligned}10 \sin^2(\theta_{12}) &= 3.18 \pm 0.16 \\ 10^2 \sin^2(\theta_{13}) &= 2.225^{+0.055}_{-0.078} \text{ (NO)} \\ &= 2.250^{+0.056}_{-0.076} \text{ (IO)} \\ 10 \sin^2(\theta_{23}) &= 5.66^{+0.16}_{-0.22} \text{ (NO)} \\ &= 5.66^{+0.18}_{-0.23} \text{ (IO)} \\ \delta/\pi &= 1.20^{+0.23}_{-0.14} \text{ (NO)} \\ &= 1.54 \pm 0.13 \text{ (IO)}\end{aligned}$$




see also: <http://globalfit.astroparticles.es>

# History of the universe




# History of the universe



## The oldest picture of the Universe



The Cosmic Microwave Background, generated at  $t \simeq 4 \times 10^5$  years

COBE (1992)    WMAP (2003)    Planck (2013)



# CMB spectra as of 2018

[Planck Collaboration, 2018]



# Big Bang Nucleosynthesis (BBN)

BBN: production of light nuclei at  $t \sim 1\text{ s}$  to  $t \sim \mathcal{O}(10^2)\text{s}$


temperature  $T_{fr} \simeq 1 \text{ MeV}$   
from nucleon freeze-out

much earlier than CMB!

strong probe for physics  
before the CMB

e.g. neutrinos!

$\nu$  affect  
universe expansion  
and  
reaction rates ( $\nu_e/\bar{\nu}_e$ )  
at BBN time...



# Big Bang Nucleosynthesis (BBN)

BBN: production of light nuclei at  $t \sim 1\text{s}$  to  $t \sim \mathcal{O}(10^2)\text{s}$


temperature  $T_{fr} \simeq 1 \text{ MeV}$   
from nucleon freeze-out

much earlier than CMB!

strong probe for physics  
before the CMB


e.g. neutrinos!

$\nu$  affect  
universe expansion  
and  
reaction rates ( $\nu_e/\bar{\nu}_e$ )  
at BBN time...




## ■ Neutrinos in the early Universe

before BBN: neutrinos coupled to plasma ( $\nu_\alpha \bar{\nu}_\alpha \leftrightarrow e^+ e^-$ ,  $\nu e \leftrightarrow \nu e$ )



## ■ Neutrinos in the early Universe

before BBN: neutrinos coupled to plasma ( $\nu_\alpha \bar{\nu}_\alpha \leftrightarrow e^+ e^-$ ,  $\nu e \leftrightarrow \nu e$ )




[Courtesy P. F. de Salas]

$T / \text{MeV}$

## ■ Neutrinos in the early Universe

before BBN: neutrinos coupled to plasma ( $\nu_\alpha \bar{\nu}_\alpha \leftrightarrow e^+ e^-$ ,  $\nu e \leftrightarrow \nu e$ )




[Courtesy P. F. de Salas]

$T / \text{MeV}$

$\nu$  decouple mostly before  $e^+ e^- \rightarrow \gamma\gamma$  annihilation!

# Neutrinos in the early Universe


before BBN: neutrinos coupled to plasma ( $\nu_\alpha \bar{\nu}_\alpha \leftrightarrow e^+ e^-$ ,  $\nu e \leftrightarrow \nu e$ )



ν decouple mostly before  $e^+ e^- \rightarrow \gamma\gamma$  annihilation!

# Neutrinos in the early Universe

before BBN: neutrinos coupled to plasma ( $\nu_\alpha \bar{\nu}_\alpha \leftrightarrow e^+ e^-$ ,  $\nu e \leftrightarrow \nu e$ )



$\nu$  decouple mostly before  $e^+ e^- \rightarrow \gamma\gamma$  annihilation!  
actually, the decoupling  $T$  is momentum dependent!

distortions to equilibrium  $f_\nu$ !

## $\nu$ oscillations in the early universe

comoving coordinates:  $a = 1/T$     $x \equiv m_e a$     $y \equiv p_a$     $z \equiv T_\gamma a$     $w \equiv T_\nu a$

density matrix:  $\varrho(x, y) = \begin{pmatrix} \varrho_{ee} \equiv f_{\nu_e} & \varrho_{e\mu} & \varrho_{e\tau} \\ \varrho_{\mu e} & \varrho_{\mu\mu} \equiv f_{\nu_\mu} & \varrho_{\mu\tau} \\ \varrho_{\tau e} & \varrho_{\tau\mu} & \varrho_{\tau\tau} \equiv f_{\nu_\tau} \end{pmatrix}$

$$\frac{d\varrho(y, x)}{dx} = \sqrt{\frac{3m_{Pl}^2}{8\pi\rho_T}} \left\{ -i \frac{x^2}{m_e^3} \left[ \frac{M_F}{2y} - \frac{2\sqrt{2}G_F y}{x^6/m_e^6} \left( \frac{E_\ell + P_\ell}{m_W^2} + \frac{4E_\nu}{3m_Z^2} \right), \varrho \right] + \frac{m_e^3 G_F^2}{(2\pi)^3 x^4 y^2} \mathcal{I}(\varrho) \right\}$$

$m_{Pl}$  Planck mass –  $\rho_T$  total energy density –  $m_{W,Z}$  mass of the  $W, Z$  bosons –  $G_F$  Fermi constant –  $[., .]$  commutator

# $\nu$ oscillations in the early universe

[Bennett, SG+, 2012.02726]

comoving coordinates:  $a = 1/T$     $x \equiv m_e a$     $y \equiv p_a$     $z \equiv T_\gamma a$     $w \equiv T_\nu a$

density matrix:  $\varrho(x, y) = \begin{pmatrix} \varrho_{ee} \equiv f_{\nu_e} & \varrho_{e\mu} & \varrho_{e\tau} \\ \varrho_{\mu e} & \varrho_{\mu\mu} \equiv f_{\nu_\mu} & \varrho_{\mu\tau} \\ \varrho_{\tau e} & \varrho_{\tau\mu} & \varrho_{\tau\tau} \equiv f_{\nu_\tau} \end{pmatrix}$

$$\frac{d\varrho(y, x)}{dx} = \sqrt{\frac{3m_{\text{Pl}}^2}{8\pi\rho_T}} \left\{ -i \frac{x^2}{m_e^3} \left[ \frac{\mathbb{M}_F}{2y} - \frac{2\sqrt{2}G_F y}{x^6/m_e^6} \left( \frac{\mathbb{E}_\ell + \mathbb{P}_\ell}{m_W^2} + \frac{4\mathbb{E}_\nu}{3m_Z^2} \right), \varrho \right] + \frac{m_e^3 G_F^2}{(2\pi)^3 x^4 y^2} \mathcal{I}(\varrho) \right\}$$

$m_{\text{Pl}}$  Planck mass –  $\rho_T$  total energy density –  $m_{W,Z}$  mass of the  $W, Z$  bosons –  $G_F$  Fermi constant –  $[\cdot, \cdot]$  commutator

$$\mathbb{M}_F = U \mathbb{M} U^\dagger$$

$$\mathbb{M} = \text{diag}(m_1^2, \dots, m_N^2)$$

$$U = R^{23} R^{13} R^{12} \quad \text{e.g. } R^{13} = \begin{pmatrix} \cos \theta_{13} & 0 & \sin \theta_{13} \\ 0 & 1 & 0 \\ -\sin \theta_{13} & 0 & \cos \theta_{13} \end{pmatrix}$$

# $\nu$ oscillations in the early universe

[Bennett, SG+, 2012.02726]

comoving coordinates:  $a = 1/T$     $x \equiv m_e a$     $y \equiv p_a$     $z \equiv T_\gamma a$     $w \equiv T_\nu a$

density matrix:  $\varrho(x, y) = \begin{pmatrix} \varrho_{ee} \equiv f_{\nu_e} & \varrho_{e\mu} & \varrho_{e\tau} \\ \varrho_{\mu e} & \varrho_{\mu\mu} \equiv f_{\nu_\mu} & \varrho_{\mu\tau} \\ \varrho_{\tau e} & \varrho_{\tau\mu} & \varrho_{\tau\tau} \equiv f_{\nu_\tau} \end{pmatrix}$

$$\frac{d\varrho(y, x)}{dx} = \sqrt{\frac{3m_{Pl}^2}{8\pi\rho_T}} \left\{ -i \frac{x^2}{m_e^3} \left[ \frac{\mathbb{M}_F}{2y} - \frac{2\sqrt{2}G_F y}{x^6/m_e^6} \left( \frac{\mathbb{E}_e + \mathbb{P}_e}{m_W^2} + \frac{4\mathbb{E}_\nu}{3m_Z^2} \right), \varrho \right] + \frac{m_e^3 G_F^2}{(2\pi)^3 x^4 y^2} \mathcal{I}(\varrho) \right\}$$

$m_{Pl}$  Planck mass –  $\rho_T$  total energy density –  $m_{W,Z}$  mass of the  $W, Z$  bosons –  $G_F$  Fermi constant –  $[\cdot, \cdot]$  commutator

$$\mathbb{M}_F = U \mathbb{M} U^\dagger$$

$$\mathbb{E}_e = \text{diag}(\rho_e, \rho_\mu, 0) \quad \mathbb{E}_\nu = S_a \left( \int dy y^3 \varrho \right) S_a \quad \text{with } S_a = \text{diag}(1, 1, 1)$$

lepton densities

neutrino densities

(only for active neutrinos)

take into account matter effects in oscillations

# $\nu$ oscillations in the early universe

[Bennett, SG+, 2012.02726]

comoving coordinates:  $a = 1/T$     $x \equiv m_e a$     $y \equiv p a$     $z \equiv T_\gamma a$     $w \equiv T_\nu a$

density matrix:  $\varrho(x, y) = \begin{pmatrix} \varrho_{ee} \equiv f_{\nu_e} & \varrho_{e\mu} & \varrho_{e\tau} \\ \varrho_{\mu e} & \varrho_{\mu\mu} \equiv f_{\nu_\mu} & \varrho_{\mu\tau} \\ \varrho_{\tau e} & \varrho_{\tau\mu} & \varrho_{\tau\tau} \equiv f_{\nu_\tau} \end{pmatrix}$

$$\frac{d\varrho(y, x)}{dx} = \sqrt{\frac{3m_{\text{Pl}}^2}{8\pi\rho_T}} \left\{ -i \frac{x^2}{m_e^3} \left[ \frac{\mathbb{M}_F}{2y} - \frac{2\sqrt{2}G_F y}{x^6/m_e^6} \left( \frac{\mathbb{E}_\ell + \mathbb{P}_\ell}{m_W^2} + \frac{4\mathbb{E}_\nu}{3m_Z^2} \right), \varrho \right] + \frac{m_e^3 G_F^2}{(2\pi)^3 x^4 y^2} \mathcal{I}(\varrho) \right\}$$

$m_{\text{Pl}}$  Planck mass –  $\rho_T$  total energy density –  $m_{W,Z}$  mass of the  $W, Z$  bosons –  $G_F$  Fermi constant –  $[\cdot, \cdot]$  commutator

$$\mathbb{M}_F = U \mathbb{M} U^\dagger \quad \mathbb{E}_\ell = \text{diag}(\rho_e, \rho_\mu, 0) \quad \mathbb{E}_\nu = S_a \left( \int dy y^3 \varrho \right) S_a$$

$\mathcal{I}(\varrho)$  collision integrals

take into account neutrino-electron scattering and pair annihilation,  
plus neutrino–neutrino interactions

2D integrals over momentum, take most of the computation time

# $\nu$ oscillations in the early universe

[Bennett, SG+, 2012.02726]

comoving coordinates:  $a = 1/T$     $x \equiv m_e a$     $y \equiv p a$     $z \equiv T_\gamma a$     $w \equiv T_\nu a$

density matrix:  $\varrho(x, y) = \begin{pmatrix} \varrho_{ee} \equiv f_{\nu_e} & \varrho_{e\mu} & \varrho_{e\tau} \\ \varrho_{\mu e} & \varrho_{\mu\mu} \equiv f_{\nu_\mu} & \varrho_{\mu\tau} \\ \varrho_{\tau e} & \varrho_{\tau\mu} & \varrho_{\tau\tau} \equiv f_{\nu_\tau} \end{pmatrix}$

$$\frac{d\varrho(y, x)}{dx} = \sqrt{\frac{3m_{Pl}^2}{8\pi\rho_T}} \left\{ -i \frac{x^2}{m_e^3} \left[ \frac{\mathbb{M}_F}{2y} - \frac{2\sqrt{2}G_F y}{x^6/m_e^6} \left( \frac{\mathbb{E}_\ell + \mathbb{P}_\ell}{m_W^2} + \frac{4\mathbb{E}_\nu}{3m_Z^2} \right), \varrho \right] + \frac{m_e^3 G_F^2}{(2\pi)^3 x^4 y^2} \mathcal{I}(\varrho) \right\}$$

$m_{Pl}$  Planck mass –  $\rho_T$  total energy density –  $m_{W,Z}$  mass of the  $W, Z$  bosons –  $G_F$  Fermi constant –  $[\cdot, \cdot]$  commutator

$$\mathbb{M}_F = U \mathbb{M} U^\dagger \quad \mathbb{E}_\ell = \text{diag}(\rho_e, \rho_\mu, 0) \quad \mathbb{E}_\nu = S_a \left( \int dy y^3 \varrho \right) S_a$$

$\mathcal{I}(\varrho)$  collision integrals

from continuity  
equation

$$\dot{\rho} = -3H(\rho + P)$$

$$\frac{dz}{dx} = \frac{\sum_{\ell=e,\mu} \left[ \frac{r_\ell^2}{r} J(r_\ell) \right] + G_1(r) - \frac{1}{2\pi^2 z^3} \int_0^\infty dy y^3 \sum_{\alpha=e}^\tau \frac{d\varrho_{\alpha\alpha}}{dx}}{\sum_{\ell=e,\mu} \left[ r_\ell^2 J(r_\ell) + Y(r_\ell) \right] + G_2(r) + \frac{2\pi^2}{15}}$$

$r = x/z$ ,  $r_\ell = m_\ell/m_e$   $r$     $J(r)$ ,  $Y(r)$  from non-relativistic transition of  $e^\pm$ ,  $\mu^\pm$   
 $G_1(r)$  and  $G_2(r)$  from electromagnetic corrections

# $\nu$ oscillations in the early universe

comoving coordinates:  $a = 1/T$     $x \equiv m_e a$     $y \equiv p a$     $z \equiv T_\gamma a$     $w \equiv T_\nu a$

density matrix:  $\varrho(x, y) = \begin{pmatrix} \varrho_{ee} \equiv f_{\nu_e} & \varrho_{e\mu} & \varrho_{e\tau} \\ \varrho_{\mu e} & \varrho_{\mu\mu} \equiv f_{\nu_\mu} & \varrho_{\mu\tau} \\ \varrho_{\tau e} & \varrho_{\tau\mu} & \varrho_{\tau\tau} \equiv f_{\nu_\tau} \end{pmatrix}$

$$\frac{d\varrho(y, x)}{dx} = \sqrt{\frac{3m_{Pl}^2}{8\pi\rho_T}} \left\{ -i \frac{x^2}{m_e^3} \left[ \frac{\mathbb{M}_F}{2y} - \frac{2\sqrt{2}G_F y}{x^6/m_e^6} \left( \frac{\mathbb{E}_\ell + \mathbb{P}_\ell}{m_W^2} + \frac{4\mathbb{E}_\nu}{3m_Z^2} \right), \varrho \right] + \frac{m_e^3 G_F^2}{(2\pi)^3 x^4 y^2} \mathcal{I}(\varrho) \right\}$$

$m_{Pl}$  Planck mass –  $\rho_T$  total energy density –  $m_{W,Z}$  mass of the  $W, Z$  bosons –  $G_F$  Fermi constant –  $[., .]$  commutator

$$\mathbb{M}_F = U \mathbb{M} U^\dagger \quad \mathbb{E}_\ell = \text{diag}(\rho_e, \rho_\mu, 0) \quad \mathbb{E}_\nu = S_a \left( \int dy y^3 \varrho \right) S_a$$

$\mathcal{I}(\varrho)$  collision integrals

from continuity equation

$$\dot{\rho} = -3H(\rho + P)$$

$$\frac{dz}{dx} = \frac{\sum_{\ell=e,\mu} \left[ \frac{r_\ell^2}{r} J(r_\ell) \right] + G_1(r) - \frac{1}{2\pi^2 z^3} \int_0^\infty dy y^3 \sum_{\alpha=e}^\tau \frac{d\varrho_{\alpha\alpha}}{dx}}{\sum_{\ell=e,\mu} \left[ r_\ell^2 J(r_\ell) + Y(r_\ell) \right] + G_2(r) + \frac{2\pi^2}{15}}$$

neutrino temperature  $w$ : same equation as  $z$ , but electrons always relativistic

# $\nu$ oscillations in the early universe

[Bennett, SG+, 2012.02726]

comoving coordinates:  $a = 1/T$     $x \equiv m_e a$     $y \equiv p a$     $z \equiv T_\gamma a$     $w \equiv T_\nu a$

density matrix:  $\varrho(x, y) = \begin{pmatrix} \varrho_{ee} \equiv f_{\nu_e} & \varrho_{e\mu} & \varrho_{e\tau} \\ \varrho_{\mu e} & \varrho_{\mu\mu} \equiv f_{\nu_\mu} & \varrho_{\mu\tau} \\ \varrho_{\tau e} & \varrho_{\tau\mu} & \varrho_{\tau\tau} \equiv f_{\nu_\tau} \end{pmatrix}$

$$\frac{d\varrho(y, x)}{dx} = \sqrt{\frac{3m_{Pl}^2}{8\pi\rho_T}} \left\{ -i \frac{x^2}{m_e^3} \left[ \frac{\mathbb{M}_F}{2y} - \frac{2\sqrt{2}G_F y}{x^6/m_e^6} \left( \frac{\mathbb{E}_\ell + \mathbb{P}_\ell}{m_W^2} + \frac{4\mathbb{E}_\nu}{3m_Z^2} \right), \varrho \right] + \frac{m_e^3 G_F^2}{(2\pi)^3 x^4 y^2} \mathcal{I}(\varrho) \right\}$$

$m_{Pl}$  Planck mass –  $\rho_T$  total energy density –  $m_{W,Z}$  mass of the  $W, Z$  bosons –  $G_F$  Fermi constant –  $[\cdot, \cdot]$  commutator

$$\mathbb{M}_F = U \mathbb{M} U^\dagger \quad \mathbb{E}_\ell = \text{diag}(\rho_e, \rho_\mu, 0) \quad \mathbb{E}_\nu = S_a \left( \int dy y^3 \varrho \right) S_a$$

$\mathcal{I}(\varrho)$  collision integrals

from continuity equation

$$\dot{\rho} = -3H(\rho + P)$$

$$\frac{dz}{dx} = \frac{\sum_{\ell=e,\mu} \left[ \frac{r_\ell^2}{r} J(r_\ell) \right] + G_1(r) - \frac{1}{2\pi^2 z^3} \int_0^\infty dy y^3 \sum_{\alpha=e}^\tau \frac{d\varrho_{\alpha\alpha}}{dx}}{\sum_{\ell=e,\mu} \left[ r_\ell^2 J(r_\ell) + Y(r_\ell) \right] + G_2(r) + \frac{2\pi^2}{15}}$$

neutrino temperature  $w$ : same equation as  $z$ , but electrons always relativistic  
initial conditions:  $\varrho_{\alpha\alpha} = \text{Fermi-Dirac at } x_{\text{in}} \simeq 0.001$ , with  $w = z \simeq 1$

# $\nu$ oscillations in the early universe

[Bennett, SG+, 2012.02726]

comoving coordinates:  $a = 1/T$     $x \equiv m_e a$     $y \equiv p a$     $z \equiv T_\gamma a$     $w \equiv T_\nu a$

density matrix:  $\varrho(x, y) = \begin{pmatrix} \varrho_{ee} \equiv f_{\nu_e} & \varrho_{e\mu} & \varrho_{e\tau} \\ \varrho_{\mu e} & \varrho_{\mu\mu} \equiv f_{\nu_\mu} & \varrho_{\mu\tau} \\ \varrho_{\tau e} & \varrho_{\tau\mu} & \varrho_{\tau\tau} \equiv f_{\nu_\tau} \end{pmatrix}$

$$\frac{d\varrho(y, x)}{dx} = \sqrt{\frac{3m_{Pl}^2}{8\pi\rho_T}} \left\{ -i \frac{x^2}{m_e^3} \left[ \frac{M_F}{2y} - \frac{2\sqrt{2}G_F y}{x^6/m_e^6} \left( \frac{E_\ell + P_\ell}{m_W^2} + \frac{4E_\nu}{3m_Z^2} \right), \varrho \right] + \frac{m_e^3 G_F^2}{(2\pi)^3 x^4 y^2} \mathcal{I}(\varrho) \right\}$$

$m_P$ : Planck mass    $\rho_T$ : total energy density    $M_F$ : mass of the  $W/Z$  bosons    $G_F$ : Fermi constant    $\mathcal{I}$ : commutator

## FORTran-Evolved Primordial Neutrino Oscillations (FortEPiaNO)

[https://bitbucket.org/ahep\\_cosmo/fortepiano](https://bitbucket.org/ahep_cosmo/fortepiano)

from continuity  
equation


$$\dot{\rho} = -3H(\rho + P)$$

will be public soon


$$\frac{dz}{dx} = \frac{\sum_{\ell=e,\mu} [r_\ell^2 J(r_\ell) + Y(r_\ell)] + G_2(r) + \frac{2\pi^2}{15}}{\frac{1}{2\pi^2 z^3} \int_0^\infty dy y^3 \sum_{\alpha=e}^\tau \frac{d\varrho_{\alpha\alpha}}{dx}}$$

neutrino temperature  $w$ : same equation as  $z$ , but electrons always relativistic  
initial conditions:  $\varrho_{\alpha\alpha} = \text{Fermi-Dirac at } x_{\text{in}} \simeq 0.001$ , with  $w = z \simeq 1$

Distortion of the momentum distribution ( $f_{\text{FD}}$ : Fermi-Dirac at equilibrium)




Distortion of the momentum distribution ( $f_{\text{FD}}$ : Fermi-Dirac at equilibrium)



$$N_{\text{eff}}^{\text{final}} = \frac{8}{7} \left( \frac{11}{4} \right)^{4/3} \frac{\rho_\nu}{\rho_\gamma} = \frac{8}{7} \left( \frac{11}{4} \right)^{4/3} \frac{1}{\rho_\gamma} \sum_i g_i \int \frac{d^3 p}{(2\pi)^3} E(p) f_{\nu,i}(p)$$

$$(11/4)^{1/3} = (T_\gamma / T_\nu)^{\text{fin}}$$


$\hookrightarrow \propto y^3 \varrho_{ii}(y)$



# Neutrino momentum distribution and $N_{\text{eff}}$


[Bennett, SG+, 2012.02726]

$$N_{\text{eff}}^{\text{any time}} = \frac{8}{7} \left( \frac{T_\gamma}{T_\nu} \right)^4 \frac{\rho_\nu}{\rho_\gamma} = \frac{8}{7} \left( \frac{T_\gamma}{T_\nu} \right)^4 \frac{1}{\rho_\gamma} \sum_i g_i \int \frac{d^3 p}{(2\pi)^3} E(p) f_{\nu,i}(p)$$



# Effect of neutrino oscillations



[Bennett, SG+, 2012.02726]





# Effect of neutrino oscillations

[Bennett, SG+, 2012.02726]

..... no  $\nu\nu$      $\sin^2\theta_{12}$      $\Delta m_{21}^2$   
- - -  $\nu\nu$ , GL     $\sin^2\theta_{13}$      $\Delta m_{31}^2$   
- - -  $\nu\nu$ , NC     $\sin^2\theta_{23}$



within  $3\sigma$  ranges allowed by global fits [deSalas, SG+, JHEP 2021]  
only  $\theta_{12}$  affects  $N_{\text{eff}}$ , at most by  $\delta N_{\text{eff}} \approx 10^{-4}$



Discretize neutrino momenta to compute integrals and evolution


two sampling methods for  $y_i$ , with  $i = 1, \dots, N_y$ :

linear spacing,

Newton-Cotes (NC) integration

Gauss-Laguerre (GL)

optimized for computing  $\int_0^\infty dy f(y)e^{-y}$



Need to define range ( $y_{\min} \leq y \leq y_{\max}$ ) and number of nodes  $N_y$

Discretize neutrino momenta to compute integrals and evolution


two sampling methods for  $y_i$ , with  $i = 1, \dots, N_y$ :

linear spacing,

Newton-Cotes (NC) integration




Gauss-Laguerre (GL)

optimized for computing  $\int_0^\infty dy f(y)e^{-y}$



Need to define range ( $y_{\min} \leq y \leq y_{\max}$ ) and number of nodes  $N_y$

Discretize neutrino momenta to compute integrals and evolution



at same  $N_y$ ,  
GL results are more stable!

GL is more efficient

$\delta N_{\text{eff}} \approx 10^{-4}$  from varying  $N_y$ ,  $y_{\max}$

## How precise is $N_{\text{eff}} = 3.04\dots$ ?

Long list of previous works... always less than  $3\nu$  mixing

[Mangano+, 2005]:  $N_{\text{eff}} = 3.046$  1st with  $3\nu$  mixing (still most cited value)

[de Salas+, 2016]:  $N_{\text{eff}} = 3.045$  updated collision terms

[SG+, 2019]:  $N_{\text{eff}} = 3.044$  more efficient and precise code,

FortEPiaNO code  $N > 3$  neutrinos allowed,  
minor differences in numerical integrals

[Bennett+, 2019]:  $N_{\text{eff}} = 3.043$  finite- $T$  QED corrections at  $\mathcal{O}(e^3)!$   
(no full calculation)

further terms should be almost negligible


[Akita+, 2020]: equations in mass and flavor basis  
 $N_{\text{eff}} = 3.044 \pm 0.0005$  approximated  $\nu\nu$  collisions

full  $\nu\nu$  interactions  
1st estimate effect of CP-violating phase

[Froustey+, 2020]:  
 $N_{\text{eff}} = 3.0440 \pm \mathcal{O}(10^{-4})$

[Bennett, SG+, 2020]:  
 $N_{\text{eff}} = 3.0440 \pm 0.0002$  1st full discussion on effect of oscillation parameters, full estimation of current numerical and physical uncertainty

FortEPiaNO improved



# $N_{\text{eff}}$ and BBN

BBN: production of light nuclei  
at  $t \sim 1\text{s}$  to  $t \sim \mathcal{O}(10^2)\text{s}$

temperature  $T_{\text{fr}} \simeq 1 \text{ MeV}$   
from nucleon freeze-out:

$$\Gamma_{n \leftrightarrow p} \sim G_F^2 T^5 = H \sim \sqrt{g_* G_N} T^2$$

$$T_{\text{fr}} \simeq (g_* G_N / G_F^4)^{1/6}$$

enters  
 $n/p = \exp(-Q/T_{\text{fr}})$

which controls element abundances

$g_*$  depends on  $N_{\text{eff}}$

abundances depend on  $N_{\text{eff}}$


$G_F$  Fermi constant

$n, p$ : neutron, proton density number

$G_N$  Newton constant

$Q = 1.293 \text{ MeV}$  neutron-proton mass difference

"(Cosmological) Relic neutrinos, from A to Z"



# $N_{\text{eff}}$ and BBN

BBN: production of light nuclei  
at  $t \sim 1\text{s}$  to  $t \sim \mathcal{O}(10^2)\text{s}$

temperature  $T_{\text{fr}} \simeq 1 \text{ MeV}$   
from nucleon freeze-out:

$$\Gamma_{n \leftrightarrow p} \sim G_F^2 T^5 = H \sim \sqrt{g_* G_N} T^2$$

$$T_{\text{fr}} \simeq (g_* G_N / G_F^4)^{1/6}$$

enters

$$n/p = \exp(-Q/T_{\text{fr}})$$

which controls element abundances

$g_*$  depends on  $N_{\text{eff}}$

abundances depend on  $N_{\text{eff}}$

$G_F$  Fermi constant


$n, p$ : neutron, proton density number

$G_N$  Newton constant

$Q = 1.293 \text{ MeV}$  neutron-proton mass difference

"(Cosmological) Relic neutrinos, from A to Z"


[Planck Collaboration, 2018]




$$N_{\text{eff}} = 2.87^{+0.24}_{-0.21}$$

(BBN only)

[Consiglio+, CPC 2018]






# B

# Bosonic neutrinos (?!? what?)

Based on:

- JCAP 03 (2018) 050



## Motivation

Neutrinos are fermions —————→ they obey Fermi-Dirac statistics

## Motivation

Neutrinos are fermions —————> they obey Fermi-Dirac statistics

Do they obey Fermi-Dirac statistics?

No experimental confirmation of spin-statistics theorem for neutrinos!

Can we find violations of the Pauli exclusion principle?

## Motivation

Neutrinos are fermions —————> they obey Fermi-Dirac statistics

Do they obey Fermi-Dirac statistics?

No experimental confirmation of spin-statistics theorem for neutrinos!

Can we find violations of the Pauli exclusion principle?

electrons

no violations for atomic electrons

e.g. look for anomalous X-rays from  
atomic decays

[Goldhaber&Scharff-Goldhaber, 1948]

[Fischbach&Kirsten&Schaeffer, 1968]

[Reines&Sobel, 1974]

...

nucleons

no violations for protons/neutrons

e.g. look for anomalous star (Sun)  
dynamics or transitions in nuclei

[Plaga, 1989]

[Miljanić+, 1990]

[Borexino, 2004]

...

see detailed discussion in [Dolgov&Smirnov, PLB 2005]

## The neutrino case

important: since spin-statistics relation confirmed for electrons,  
difficult to imagine large deviation for neutrinos

## The neutrino case

important: since spin-statistics relation confirmed for electrons,  
difficult to imagine large deviation for neutrinos

violation of the Pauli principle for  $\nu$  should show up  
in elementary processes where identical  $\nu$  are involved

for example the two-neutrino double beta decay,



## The neutrino case

important: since spin-statistics relation confirmed for electrons,  
difficult to imagine large deviation for neutrinos

violation of the Pauli principle for  $\nu$  should show up  
in elementary processes where identical  $\nu$  are involved

for example the two-neutrino double beta decay,  
 $A \rightarrow A' + 2\bar{\nu} + 2e^-$  or  $A \rightarrow A' + 2\nu + 2e^+$

Fermi-Bose parameter  $\kappa_\nu$  [Dolgov+, JCAP 2005]

$$f_\nu(E) = \frac{1}{\exp(E/T) + \kappa_\nu}$$

“mixed”  
distribution!

$$\text{BE} \leftarrow \kappa_\nu = -1 \xleftarrow[\text{MB}]{\kappa_\nu = 0} \xrightarrow{\kappa_\nu = +1} \text{FD}$$

[Barabash+, NPB 2007]:  $\kappa_\nu \gtrsim -0.2$

100% violation excluded [Barabash+, NPB 2007],  
but still 50% admixture of bosonic component allowed

## Constraints on $\kappa_\nu$ from BBN

[de Salas, SG+, JCAP 03 (2018) 050]

what can cosmology say about  $\kappa_\nu$ ?

different  $f_\nu(p)$  affects BBN!

statistics factor becomes  $(1 - \kappa_\nu f_\nu)$

$(1 + f_\nu) \rightarrow$  Bose enhancement,

$(1 - f_\nu) \rightarrow$  Pauli blocking

# Constraints on $\kappa_\nu$ from BBN

[de Salas, SG+, JCAP 03 (2018) 050]

what can cosmology say about  $\kappa_\nu$ ?

different  $f_\nu(p)$  affects BBN!

statistics factor becomes  $(1 - \kappa_\nu f_\nu)$

$(1 + f_\nu) \rightarrow$  Bose enhancement,


$(1 - f_\nu) \rightarrow$  Pauli blocking



change of n/p ratio at BBN

[Dolgov+, JCAP 2005]

less He, more D, less Li



deviation from  $\kappa_\nu = 1$   
obtained with a modified version  
of PArthENoPE  
[Consiglio+, CPC 2018]

# Constraints on $\kappa_\nu$ from BBN

[de Salas, SG+, JCAP 03 (2018) 050]

what can cosmology say about  $\kappa_\nu$ ?

different  $f_\nu(p)$  affects BBN!

statistics factor becomes  $(1 - \kappa_\nu f_\nu)$

$(1 + f_\nu) \rightarrow$  Bose enhancement,

$(1 - f_\nu) \rightarrow$  Pauli blocking



change of n/p ratio at BBN



[Dolgov+, JCAP 2005]

less He, more D, less Li

He or D alone cannot constrain  $\kappa_\nu$

Li problem drives  $\omega_b$  down  
and  $\kappa_\nu$  to -1

also when prior on  $\omega_b$  is included



# Neutrino densities and $\kappa_\nu$

$$f_\nu(E) = \frac{1}{\exp(E/T) + \kappa_\nu}$$

$\kappa_\nu$  affects  
background evolution:

$$\rho_\nu^{\text{rel}} \simeq \frac{g_\nu}{2\pi^2} \int_0^\infty dp p^3 f_\nu(p)$$

bosons:

$$\frac{\pi^2}{30} g_i T^4$$

$$\rho_\nu^{\text{nr}} \simeq m_\nu \frac{g_\nu}{2\pi^2} \int_0^\infty dp p^2 f_\nu(p)$$

bosons:

$$\frac{\zeta(3)}{\pi^2} m_\nu g_i T^3$$


fermions:

$$\frac{7}{8} \frac{\pi^2}{30} g_i T^4$$

$$\frac{3}{4} \frac{\zeta(3)}{\pi^2} m_\nu g_i T^3$$

changing  $\kappa_\nu$  "mimics" altering  $N_{\text{eff}}$  or  $\sum m_\nu$  (at late or early times)

partial degeneracies with  $N_{\text{eff}}$  and  $\sum m_\nu$



## CMB/BAO constraints on $\kappa_\nu$

need to cover  $\kappa_\nu - \sum m_\nu$  degeneracy:  
vary both!

degeneracy affects  
mostly CMB only bounds

with BAO, bound on  $\sum m_\nu$  is stronger

adding radiation (through  $\kappa_\nu$ ) and  $\Omega_\Lambda$  alters  
 $H_0$  and compensates a bit the larger mass



bounds:  $\kappa_\nu \gtrsim -0.1$  at 68%

$-1 \leq \kappa_\nu \leq 1$  at 95%

$\kappa_\nu = -1$  corresponds to  
 $N_{\text{eff}} \simeq 3.47$  at early times

inside Planck  $2\sigma$  region!  
reasonably it's not excluded

[de Salas, SG+, JCAP 03 (2018) 050]




## C

# Clustering in the local Universe

Based on:

- JCAP 09 (2017) 034
- JCAP 01 (2020) 015



Relic neutrinos are slow!  $[c_\nu \sim 160(1+z)(1\text{ eV}/m_\nu) \text{ km s}^{-1}]$

Can be trapped in the gravitational potential of the Milky Way and neighbours

$f_c(m_i) = n_i/n_{i,0}$  clustering factor → How to compute it?

Idea from [Ringwald & Wong, 2004] → **N-one-body** =  $N \times$  single  $\nu$  simulations

→ each  $\nu$  evolved from initial conditions at  $z = 3$

→ spherical symmetry, coordinates  $(r, \theta, p_r, l)$

→ need  $\rho_{\text{matter}}(z) = \rho_{\text{DM}}(z) + \rho_{\text{baryon}}(z)$

### Assumptions:

$\nu$ s are independent

only gravitational interactions

$\nu$ s do not influence matter evolution

$$(\rho_\nu \ll \rho_{\text{DM}})$$

how many  $\nu$ s is "N"?

→ must sample all possible  $r, p_r, l$

→ must include all possible  $\nu$ s that reach the MW


(fastest ones may come from several (up to  $\mathcal{O}(100)$ ) Mpc!)

given  $N \nu$ :

→ weigh each neutrinos

## Forward-tracking and back-tracking


initial phase space,  $z = 4 \longrightarrow$  homogeneous Fermi-Dirac distribution


final phase space,  $z = 0$

## Forward-tracking and back-tracking

initial phase space,  $z = 4 \longrightarrow$  homogeneous Fermi-Dirac distribution




compute final position of each particle




final phase space,  $z = 0$

## Forward-tracking and back-tracking

initial phase space,  $z = 4 \longrightarrow$  homogeneous Fermi-Dirac distribution




compute final position of each particle




final phase space,  $z = 0$

## Forward-tracking and back-tracking

initial phase space,  $z = 4 \longrightarrow$  homogeneous Fermi-Dirac distribution




use positions to find neutrino distribution today




final phase space,  $z = 0$

## Forward-tracking and back-tracking

initial phase space,  $z = 4 \longrightarrow$  homogeneous Fermi-Dirac distribution



only interested in overdensity at Earth? ★




a lot of time is wasted!


final phase space,  $z = 0$

## Forward-tracking and back-tracking

initial phase space,  $z = 4 \longrightarrow$  homogeneous Fermi-Dirac distribution



only interested in overdensity at Earth? ★



a lot of time is wasted!

smarter way: track backwards  
only interesting particles!

final phase space,  $z = 0$

## Advantages of tracking back

First advantage is in computational terms: much less points to compute

## Advantages of tracking back

First advantage is in computational terms: much less points to compute

Second advantage: no need to use spherical symmetry!

### Forward-tracking

initial conditions need to sample  
1D for position + 2D for momentum  
when using spherical symmetry

with full grid would require 3+3 dimensions!

Impossible to relax spherical symmetry!

### Back-tracking


"Initial" conditions only described by 3D in momentum  
(position is fixed, apart for checks)

can do the calculation with any astrophysical setup

## Advantages of tracking back


First advantage is in computational terms: much less points to compute

Second advantage: no need to use spherical symmetry!




# Clustering results with back-tracking

In comparison with previous results:




# Clustering results with back-tracking

In comparison with previous results:



# Clustering results with back-tracking

In comparison with previous results:



**Warning:** NFW  
is not the same  
for all the cases!

[de Salas+, 2017]

and

[Zhang<sup>2</sup>, 2018]


use  $\gamma \neq 1$ ,  
now we have

$$\gamma = 1$$

[Ringwald&Wong,  
2004] uses old  
parameters

# Clustering results with back-tracking

In comparison with previous results:



**Warning:** NFW  
is not the same  
for all the cases!

[de Salas+, 2017]

and

[Zhang<sup>2</sup>, 2018]

use  $\gamma \neq 1$ ,  
now we have

$$\gamma = 1$$

[Ringwald&Wong,  
2004] uses old  
parameters


# D

# Direct Detection

i.e. currently science-fiction, but in few years...

Based on:


- arxiv:1808.01892
- JCAP 07 (2019) 047



## The oldest picture of the Universe

The Cosmic Microwave Background, generated at  $t \simeq 4 \times 10^5$  years


COBE (1992)    WMAP (2003)    Planck (2013)



## The oldest picture of the Universe

The Cosmic Neutrino Background, generated at  $t \simeq 1$  s

$\dots \rightarrow 2019 \rightarrow \dots$




## How to directly detect non-relativistic neutrinos?

Remember that  $\langle E_\nu \rangle \simeq \mathcal{O}(10^{-4})$  eV today → a process without energy threshold is necessary

[Weinberg, 1962]: neutrino capture in  $\beta$ -decaying nuclei  $\nu + n \rightarrow p + e^- + \bar{\nu}$

Main background:  $\beta$  decay  $n \rightarrow p + e^- + \bar{\nu}$ !

signal is a peak at  $2m_\nu$   
above  $\beta$ -decay endpoint  
only with a lot of material  
need a very good energy resolution



$$\frac{d\tilde{\Gamma}_{\text{CNB}}}{dE_e}(E_e) = \frac{1}{\sqrt{2\pi}\sigma} \sum_{i=1}^{N_\nu} \bar{\sigma} N_T |U_{ei}|^2 n_0 f_c(m_i) \times e^{-\frac{[E_e - (E_{\text{end}} + m_i + m_{\text{lightest}})]^2}{2\sigma^2}}$$

$$\frac{d\Gamma_\beta}{dE_e} = \frac{\bar{\sigma}}{\pi^2} N_T \sum_{i=1}^{N_\nu} |U_{ei}|^2 H(E_e, m_i)$$

$$\frac{d\tilde{\Gamma}_\beta}{dE_e}(E_e) = \frac{1}{\sqrt{2\pi}\sigma} \int_{-\infty}^{+\infty} dx \frac{d\Gamma_\beta}{dE_e}(x) \exp\left[-\frac{(E_e - x)^2}{2\sigma^2}\right]$$



# $\beta$ and Neutrino Capture spectra

[PTOLEMY, JCAP 07 (2019) 047]

$$\frac{d\tilde{\Gamma}_{\text{CNB}}}{dE_e}(E_e) = \frac{1}{\sqrt{2\pi}\sigma} \sum_{i=1}^{N_\nu} \bar{\sigma} N_T |U_{ei}|^2 n_0 f_c(m_i) \times e^{-\frac{[E_e - (E_{\text{end}} + m_i + m_{\text{lightest}})]^2}{2\sigma^2}}$$

$$\frac{d\Gamma_\beta}{dE_e} = \frac{\bar{\sigma}}{\pi^2} N_T \sum_{i=1}^{N_\nu} |U_{ei}|^2 H(E_e, m_i)$$

$$\frac{d\tilde{\Gamma}_\beta}{dE_e}(E_e) = \frac{1}{\sqrt{2\pi}\sigma} \int_{-\infty}^{+\infty} dx \frac{d\Gamma_\beta}{dE_e}(x) \exp\left[-\frac{(E_e - x)^2}{2\sigma^2}\right]$$



Pontecorvo Tritium Observatory for Light, Early-universe, Massive-neutrino Yield (PTOLEMY)

expected resolution  $\Delta \simeq 0.1 \text{ eV}$ ?  
 $0.05 \text{ eV}?$

can probe  $m_\nu \simeq 1.4\Delta \simeq 0.1 \text{ eV}$

built mainly for CNB

$M_T = 100 \text{ g of atomic } {}^3\text{H}$

$$\Gamma_{\text{CNB}} = \sum_{i=1}^3 |U_{ei}|^2 [n_i(\nu_{h_R}) + n_i(\nu_{h_L})] N_T \bar{\sigma}$$

$\sim \mathcal{O}(10) \text{ yr}^{-1}$

$N_T$  number of  ${}^3\text{H}$  nuclei in a sample of mass  $M_T$        $\bar{\sigma} \simeq 3.834 \times 10^{-45} \text{ cm}^2$        $n_i$  number density of neutrino  $i$

(without clustering)

Pontecorvo Tritium Observatory for Light, Early-universe, Massive-neutrino Yield (PTOLEMY)

expected resolution  $\Delta \simeq 0.1 \text{ eV}$ ?  
 $0.05 \text{ eV}?$

can probe  $m_\nu \simeq 1.4\Delta \simeq 0.1 \text{ eV}$

built mainly for CNB

$M_T = 100 \text{ g of atomic } {}^3\text{H}$

enhancement from  
 $\nu$  clustering in the galaxy?

enhancement from  
other effects?

$$\Gamma_{\text{CNB}} = \sum_{i=1}^3 |U_{ei}|^2 [\textcolor{red}{n}_i(\nu_{h_R}) + \textcolor{red}{n}_i(\nu_{h_L})] N_T \bar{\sigma}$$

$N_T$  number of  ${}^3\text{H}$  nuclei in a sample of mass  $M_T$        $\bar{\sigma} \simeq 3.834 \times 10^{-45} \text{ cm}^2$        $n_i$  number density of neutrino  $i$

(without clustering)

$\sim \mathcal{O}(10) \text{ yr}^{-1}$



using the definition:

$$N_{\text{th}}^i(\theta) = A_\beta N_\beta^i(\hat{E}_{\text{end}} + \Delta E_{\text{end}}, m_i, U) + \mathbf{A}_{\text{CNB}} N_{\text{CNB}}^i(\hat{E}_{\text{end}} + \Delta E_{\text{end}}, m_i, U) + N_b$$

if  $\mathbf{A}_{\text{CNB}} > 0$  at  $N\sigma$ , direct detection of CNB accomplished at  $N\sigma$

statistical only!

significance on  $A_{\text{CNB}} > 0$   
as a function of  $\hat{m}_{\text{lightest}}$ ,  $\Delta$



E-R

(skipping...)

seriously, I cannot go  
through the entire alphabet in 50 minutes!


## S

## (Light) Sterile neutrinos

let's pretend they exist

Based on:

- JPG 43 (2016) 033001
- JHEP 06 (2017) 135
- PLB 782 (2018) 13-21
- in preparation
- JCAP 07 (2019) 014
- arxiv:2003.02289
- JCAP 07 (2019) 047



Problem: anomalies  
in SBL experiments

→ { errors in flux calculations?  
deviations from 3- $\nu$  description?

A short review:

**LSND** search for  $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ , with  $L/E = 0.4 \div 1.5$  m/MeV. Observed a  $3.8\sigma$  excess of  $\bar{\nu}_e$  events [Aguilar et al., 2001]

**Reactor** re-evaluation of the expected anti-neutrino flux  $\Rightarrow$  disappearance of  $\bar{\nu}_e$  events compared to predictions ( $\sim 3\sigma$ ) with  $L < 100$  m  
[Mention et al, 2011], [Azabajan et al, 2012]

**Gallium** calibration of GALLEX and SAGE Gallium solar neutrino experiments give a  $2.7\sigma$  anomaly (disappearance of  $\nu_e$ )  
[Giunti, Laveder, 2011]

MiniBooNE


See next

Possible explanation:


Additional squared mass difference  $\Delta m_{\text{SBL}}^2 \simeq 1 \text{ eV}^2$

# $\nu_s$ at reactors in 2020


[RENO+NEOS, 2020]




[DANSS, 2020]




[Neutrino-4, PZETF 2020]




[PROSPECT, PRD 2020]




[STEREO, PRD 2020]




[SoLiD, JINST 2018]






claimed  $> 3\sigma$   
preference for  
3+1 over 3ν case


best fit  
incompatible  
with other  
reactor  
experiments



energy resolution smearing not properly taken into account?



proper energy resolution treatment  
moves best-fit  $\rightarrow \sin^2 2\vartheta \simeq 1$






need to take into account  
violation of Wilk's theorem

relaxed constraints

purpose: check LSND signal

$L \simeq 541$  m,  $200$  MeV  $\leq E \lesssim 3$  GeV




no money, no near detector





purpose: check LSND signal

$$L \simeq 541 \text{ m}, 200 \text{ MeV} \leq E \lesssim 3 \text{ GeV}$$

no money, no near detector






first indication in favor of sterile from  $\nu_\mu$  DIS!

although rather weak:  $\log_{10} BF \simeq 1$  (weak preference)  
 or compatible with no oscillations at  $p$ -value of 8%

## $N_{\text{eff}}$ and the new mixing parameters

[SG+, JCAP 07 (2019) 014]


Only vary one angle and fix two to zero: do they have the same effect?



## $N_{\text{eff}}$ and the new mixing parameters

[SG+, JCAP 07 (2019) 014]



We can vary more than one angle:



## $N_{\text{eff}}$ and the new mixing parameters

[SG+, JCAP 07 (2019) 014]



We can vary more than one angle:



## $N_{\text{eff}}$ and the new mixing parameters



[SG+, JCAP 07 (2019) 014]

We can vary more than one angle:




Cosmological constraints are stronger than most other probes

But much more model dependent (as all the cosmological constraints)!



Cosmological constraints are stronger than most other probes

But much more model dependent (as all the cosmological constraints)!





Warning: tension between reactor experiments and CMB bounds!

# Can new physics solve the anomalies and tensions?

Many attempts to explain LSND/MiniBooNE anomalies,  
APP vs DIS, oscillations vs cosmo tensions with new physics

one recent example: [Dentler+, 2019]

$\mathcal{L} \supset -g\bar{\nu}_s\nu_s\phi$  with  $\mathcal{O}(\text{eV}) \lesssim m_4 \lesssim \mathcal{O}(100 \text{ keV})$  and  $m_\phi \lesssim m_4$   
↳ new interactions with scalar  $\phi$  and  $\nu_s$  decay





see also: [de Gouvea+, 2019], [Moulai+, 2019], [Fischer+, 2019], [Diaz+, 2019], ...

# Can new physics solve the anomalies and tensions?

Many attempts to explain LSND/MiniBooNE anomalies,  
APP vs DIS, oscillations vs cosmo tensions with new physics

one recent example: [Dentler+, 2019]

$\mathcal{L} \supset -g\bar{\nu}_s\nu_s\phi$  with  $\mathcal{O}(\text{eV}) \lesssim m_4 \lesssim \mathcal{O}(100 \text{ keV})$  and  $m_\phi \lesssim m_4$   
↳ new interactions with scalar  $\phi$  and  $\nu_s$  decay



see also: [de Gouvea+, 2019], [Moultai+, 2019], [Fischer+, 2019], [Diaz+, 2019], ...

## Can new physics solve the anomalies and tensions?

Many attempts to explain LSND/MiniBooNE anomalies,  
APP vs DIS, oscillations vs cosmo tensions with new physics

another example: [Liao+, 2019]

$$\mathcal{L}_{\text{NC-NSI}} = -2\sqrt{2} G_F \epsilon_{\alpha\beta}^{fC} [\bar{\nu}_\alpha \gamma^\rho P_L \nu_\beta] [\bar{f} \gamma_\rho P_C f]$$

$$\mathcal{L}_{\text{CC-NSI}} = -2\sqrt{2} G_F \epsilon_{\alpha\beta}^{ff' C} [\bar{\nu}_\beta \gamma^\rho P_L \ell_\alpha] [\bar{f}' \gamma_\rho P_C f]$$

Non-standard interactions (NSI) involving  $\nu_s$

# Can new physics solve the anomalies and tensions?

Many attempts to explain LSND/MiniBooNE anomalies,  
APP vs DIS, oscillations vs cosmo tensions with new physics


another example: [Liao+, 2019]

$$\mathcal{L}_{\text{NC-NSI}} = -2\sqrt{2}G_F \epsilon_{\alpha\beta}^{fc} [\bar{\nu}_\alpha \gamma^\rho P_L \nu_\beta] [\bar{f} \gamma_\rho P_C f]$$


$$\mathcal{L}_{\text{CC-NSI}} = -2\sqrt{2}G_F \epsilon_{\alpha\beta}^{ff'c} [\bar{\nu}_\beta \gamma^\rho P_L \ell_\alpha] [\bar{f}' \gamma_\rho P_C f]$$

Non-standard interactions (NSI) involving  $\nu_s$

MINOS+  
vs APP



IceCube/  
DeepCore  
vs APP








# Conclusions

almost there!





# What do we learn from relic neutrinos?

D






Direct detection - wonderful opportunities for the future



I Indirect probes - what we have now, it's a lot and it will improve






N New physics - beyond the corner? neutrinos will help us find it!







# What do we learn from relic neutrinos?

D






Direct detection - wonderful opportunities for the future



I Indirect probes - what we have now, it's a lot and it will improve



N New physics - beyond the corner? neutrinos will help us find it!



Thank you for the attention!